

4-10-2000

## Soybean seed treatments for 2000

X. B. Yang

*Iowa State University*, [xbyang@iastate.edu](mailto:xbyang@iastate.edu)

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Plant Pathology Commons](#)

---

### Recommended Citation

Yang, X. B., "Soybean seed treatments for 2000" (2000). *Integrated Crop Management News*. 2084.  
<http://lib.dr.iastate.edu/cropnews/2084>

**The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.**

---

# Soybean seed treatments for 2000

## **Abstract**

Dry weather conditions are predicted for this year's growing season. Although soybean seedling disease is generally not a concern under dry conditions, I have received questions on

1. seed treatments because of the early planting of soybeans in March
2. the effects of seed discoloration on seed quality, and
3. new seed treatment chemicals on the market.

## **Keywords**

Plant Pathology

## **Disciplines**

Agricultural Science | Agriculture | Plant Pathology



## Soybean seed treatments for 2000

Dry weather conditions are predicted for this year's growing season. Although soybean seedling disease is generally not a concern under dry conditions, I have received questions on

1. seed treatments because of the early planting of soybeans in March
2. the effects of seed discoloration on seed quality, and
3. new seed treatment chemicals on the market.

### Early planting

In a normal-weather season, early-planted soybeans would grow in low-temperature, high-moisture soil, which increases seedling disease risk. If this spring's weather conditions are dry, seedling disease risk is less of a concern because dry soil is not conducive for disease occurrence. However, if soybean seeds are planted in fields with high moisture, such as river bottom fields, seed treatments may be needed. In fields with high moisture, disease risk may be high for early-planted soybean. Cool ( $<60^{\circ}\text{F}$ ) and wet soils are conducive to fungal seed rot or seedling death caused by *Pythium* and *Fusarium*.



[1] Damping-off caused by *Pythium*.

A study at Iowa State University (ISU) showed that most problematic *Pythium* species in Iowa prefer cool soil temperatures ( $<60^{\circ}\text{F}$ ) for infection.

### Seed quality

Severe seed discoloration is an indicator of potential low-quality soybean seeds. However, only some causes of discoloration are associated with reduced germination rate. Discoloration may be caused by stress, insect wounding, or disease. It is unlikely that stress reduces seed quality, and thus germination, as much as viral or fungal diseases. Severe infection by viral and fungal diseases may result in reduced seed germination and is associated with seed discoloration. Seed treatment for seeds infected by fungal pathogens can improve germination rate. Treatments for seed infected with viruses are not available.

### New chemical treatments

Chemicals are effective in controlling some fungi and these chemicals are often specific for certain diseases. Only the use of the correct chemicals can prevent damping-off. Refer to previous ICM articles on seed and seedling treatments for lists of chemicals and their control efficacies for major soybean seedling pathogens in Iowa. Generally, Apron is effective for controlling *Pythium* and *Phytophthora*.

ApronMaxx is a new soybean seed treatment. It is a prepack formulation of Maxim (fludioxonil at 2.5 g/100 kg seed) and Apron XL (at 3.75g/100 kg seed). This seed treatment chemical can effectively control a broad spectrum of soybean seedling diseases, including almost all important seedling diseases in Iowa. It reduces fungal damage by contact and systematic control. Field evaluations by scientists at the Universities of Illinois, Missouri, and Wisconsin as well as ISU show an increase in yield and stand establishment. This chemical also is very effective against *Phomopsis*, a major fungus causing seed-quality problems in the past 2 years. Furthermore, this treatment can be used for on-farm seed treatments with simple equipment mounted directly and conveniently on a wagon or a truck box to dispense fungicides onto seeds during planting.

This article originally appeared on page 27 of the IC-484 (4) -- April 10, 2000 issue.

---

**Source URL:**

<http://www.ipm.iastate.edu/ipm/icm//ipm/icm/2000/4-10-2000/soytreat2000.html>

**Links:**

[1] <http://www.ipm.iastate.edu/ipm/icm//idamp.html>

IOWA STATE UNIVERSITY  
University Extension